

¡ Important information, read carefully !

**Syrup**

**Transpulmin®**  
**Antitussive and secretolytic**

**Composition:**

5 mL contain: Pipazethate HCl 10 mg (Selvigon®), Guaifenesin 25 mg, Saccharose 1.75 g, Glucose 1.75 g, Sorbitol 0.35 g (equivalent to 3.85 g Glucose), Sodium Saccharin, and other excipients q.s.

**Indications:**

All types of irritant and convulsant cough of every genesis and degree of severity. Cough with inflammatory and non inflammatory diseases of the respiratory airways.

**Contraindications:**

In case of hypersensitivity toward these substances. Although so far there have been no signs indicating an adverse effect of **Transpulmin®** during pregnancy. Pregnant women are requested to inform their physician about the ingestion of this drug. This precaution serves for necessary control on part of the physician of all drugs taken during pregnancy.

**Warnings and Cautions:**

May alter the reactivity.

For diabetic patients, 10 mL contain 7.7 g of glucose.

If discomforts persist for more than 3 days, worsen or new discomforts appear, ask your physician.

Do not exceed the recommended dose.

**Dosage:**

Unless otherwise prescribed :

Adults: 2 teaspoonfuls 3-4 times daily, as required.

Children over 3 years: 1 teaspoonful 3-4 times daily.

Infants (1 -3 years): 1/2 teaspoonful 3-4 times daily.

Maximum dose of treatment:

Adults: Do not exceed 80 mL daily.

Children over 3 years: Do not exceed 40 mL daily.

Infants (1 -3 years): Do not exceed 20 mL daily.

Maximum duration of treatment: 10 days.

**Storage:**

Do not use after the expiration date on the package.

Do not store above 30° C.

Do not use longer than 1 year after bottle was first opened.

**KEEP DRUGS OUT OF REACH OF CHILDREN.**

**Presentation:**

**Transpulmin® Syrup:** bottle containing 100 mL.